

Strongly Correlated States of Ultracold Rotating Dipolar Fermi Gases

Klaus Osterloh¹, Nuria Barberán², and Maciej Lewenstein³

(1) Institute for Theoretical Physics, University of Hannover, Appelstr. 2, D-30167 Hannover, Germany

(2) Dept. ECM, Facultat de Física, Universitat de Barcelona, E-08028 Barcelona, Spain and

(3) ICFO - Institut de Ciències Fotòniques, Parc Mediterrani de la Tecnologia, Spain

(Dated: February 6, 2008)

We study strongly correlated ground and excited states of rotating quasi-2D Fermi gases constituted of a small number of dipole-dipole interacting particles with dipole moments polarized perpendicular to the plane of motion. As the number of atoms grows, the system enters an *intermediate regime*, where ground states are subject to a competition between distinct bulk-edge configurations. This effect obscures their description in terms of composite fermions and leads to the appearance of novel composite fermion quasi-hole states. In the presence of dipolar interactions, the principal Laughlin state at filling $\nu = 1/3$ exhibits a substantial energy gap for neutral (total angular momentum conserving) excitations, and is well-described as an incompressible Fermi liquid. Instead, at lower fillings, the ground state structure favors crystalline order.

PACS numbers: 03.75.Ss, 73.43.-f

Some of the most fascinating challenges of modern atomic and molecular physics arguably concern ultracold dipolar quantum gases [1]. The recent experimental realization of a quantum degenerate dipolar Bose gas of Chromium [2], and the progress in trapping and cooling of dipolar molecules [3] have opened the path towards ultracold quantum gases with dominant dipole interactions. Particularly interesting in this context are *rotating dipolar gases* (RDG). Bose-Einstein condensates of RDGs exhibit novel forms of vortex lattices, e. g., square, stripe- and bubble-“crystal” lattices [4]. The stability of these phases in the lowest Landau level was recently investigated [5]. We have demonstrated that the quasi-hole gap survives the large N limit for fermionic RDGs [6]. This property makes them perfect candidates to approach the strongly correlated regime, and to realize Laughlin liquids (cf. [7]) at filling $\nu = 1/3$, and quantum Wigner crystals at $\nu \leq 1/7$ [8] for a mesoscopic number of atoms $N \simeq 50 - 100$. Lately, Rezayi et al. [9] have shown that the presence of a small amount of dipole-dipole interactions stabilizes the so-called bosonic Rezayi-Read state at $\nu = 3/2$ whose excitations are both fractional and non-Abelian.

In this letter, we investigate quasi-2D microscopic and mesoscopic clouds of fermions whose dipole moments are polarized perpendicular to their plane of motion. Results are obtained by exact diagonalization of the Hamiltonian of the system for $N = 3 - 12$ particles. We study the lowest Landau level (LLL) physics, in particular the crossover from the weakly interacting Fermi liquid regime to strongly correlated fractional quantum Hall-like states. As N grows, the ground state particles either organize themselves in the bulk or crystallize on the edge. This competition is based on inter-particle correlations and exceeds the scope of the effective theory of composite fermions [10]. It leads to the appearance of novel ground states with a quasi-hole at the origin which are linked to unfavored composite-fermion states. The analysis of the principal Laughlin state at filling $\nu = 1/3$ reveals that its low energy excitations in the angular momentum subspaces where quasi-holes are to be found, correspond to edge excitations

rather than topological quasi-hole defects. Instead, “neutral” particle-hole pair excitations are already substantially gapped in these microscopic systems.

We consider a system of N dipolar fermions rotating in an axially symmetric harmonic trapping potential strongly confined in the direction of the axis of rotation. Along this z -axis, the dipole moments, as well as spins are assumed to be aligned. Various ways of experimental realizations of ultracold dipolar gases are discussed in [1]. In case of low temperature T and weak chemical potential μ with respect to the axial confinement ω_z , the gas is effectively two-dimensional, and the Hamiltonian of the system in the rotating reference frame reads

$$\mathcal{H} = \sum_{j=1}^N \frac{1}{2M} (\vec{p}_j - M\Omega\vec{e}_z \times \vec{r}_j)^2 + \frac{M}{2} (\omega_0^2 - \Omega^2) r_j^2 + V_d(1)$$


Here, $\omega_0 \ll \omega_z$ is the radial trap frequency, Ω is the frequency of rotation, M is the mass of the particles, $V_d = \sum_{j < k}^N \frac{d^2}{|\vec{r}_j - \vec{r}_k|^3}$ is the dipolar interaction potential (rotationally invariant with respect to the z -axis), d is the dipole moment, and $\vec{r}_j = x_j\vec{e}_x + y_j\vec{e}_y$ is the position vector of the j -th particle. The first term of (1) is formally equivalent to the Landau Hamiltonian of particles with mass M and charge e moving in a constant magnetic field of strength $B = 2M\Omega c/e$ perpendicular to their plane of motion. The spectrum of $\mathcal{H}_{\text{Landau}}$ consists of equidistantly spaced and highly degenerate Landau levels (LL) with energies $\varepsilon_n = \hbar\omega_c(n + 1/2)$ where $\omega_c = 2\Omega$. We denote by $N_{\text{LL}} = 1/2\pi l^2$ the number of states per unit area in each LL, where $l = \sqrt{\hbar/m\omega_c}$ is the magnetic length. Given a homogeneous fermionic surface density n_f , the filling factor $\nu = 2\pi l^2 n_f$ can be defined and denotes the fraction of occupied Landau levels. Even though the above definition applies to infinite homogeneous systems, it may be used for finite systems as a suitable truncation of the Hilbert space at specific angular momenta. The second term in (1) accounts for a rotationally induced effective reduction of the trap strength. For critical rotation $\Omega \rightarrow \omega_0$, the confining potential vanishes.

In the following, it is assumed that particles solely occupy the LLL [11]. Restricted to the LLL and rewritten in second quantized form, the Hamiltonian (1) reads

$$\hat{\mathcal{H}} = \hbar\omega_0\hat{N} + \hbar(\omega_0 - \Omega)\hat{L}^z + \sum_{m_1, \dots, m_4} V_{1234} a_{m_1}^\dagger a_{m_2}^\dagger a_{m_4} a_{m_3} \quad (2)$$

where $\hat{N}$ and $\hat{L}^z$ are the total number and z-component angular momentum operators, $a_{m_i}^\dagger$ creates a particle with angular momentum m_i , and $V_{1234} = \frac{1}{2} \langle m_1 m_2 | V_d | m_3 m_4 \rangle$ is the matrix element of interaction expressed in the Fock–Darwin single particle angular momentum basis [12]. Due to circular symmetry, (2) can be diagonalized blockwise for a given L^z . Calculations have been performed for $N = 3$ to 12 particles with complete and Davidson block diagonalization techniques [13], respectively. In the LLL Hamiltonian, besides the first constant term, there are two competing contributions to the energy: the kinetic term linear in $\hat{L}$ and the strength of interaction, which scales as $d^2/l^3 = 2\hbar\omega_0(a_d/l)$ with $a_d = Md^2/\hbar^2$. The natural unit of energy is $\hbar\omega_0$, whereas distances are measured in l ; from now on, we set $a_d/l \equiv 1$. We analyze the ground state interaction energy for a given L^z . It reveals plateaus, clearly visible for small N (see Fig. 1).

FIG. 1: Interaction contribution to the GS energy as a function of L^z




Ground state candidates are the first states of these plateaus where a downward cusp occurs in the spectrum. By tuning the rotational frequency, some of these states are selected as true ground states at specific “magic” angular momenta as depicted in Fig. 2. For relatively low values of Ω , the ground state is the filled LL state at $\nu = 1$, which is completely insensitive to type and strength of interaction as long as the LLL approximation holds.

When Ω is continuously increased, the system evolves from the weakly interacting regime (where the second term of Eq.(2) is larger than the third) to strongly correlated states (where the interaction is dominant). In case of short-range interactions, this process terminates at $L_0^z = \nu_0^{-1}N(N-1)/2$, where the fermionic (bosonic) Laughlin states at filling $\nu_0 = 1/3(1/2)$ become the true ground states.

The existence of a final L^z is due to the fact that the ground state contact interaction energy vanishes for $L^z \geq L_0^z$.

FIG. 2: Ground state angular momentum series over $\alpha \equiv \omega_0 - \Omega$ (the divergence at $\alpha = 0$ is not shown).




The long range nature of dipolar interactions lifts this degeneracy. Thus, the whole principal series of fillings, i.e., $\nu = 1/(2m+1)$ for fermionic gases, is accessible. To reveal the internal structures of relevant states, we consider the density-density correlation function $\hat{\rho}(\vec{r}, \vec{r}_0)$, which represents the conditional probability to find one atom at $\vec{r}$ when another is simultaneously located at $\vec{r}_0$,

$$\hat{\rho}(\vec{r}, \vec{r}_0) = \sum_{j < k}^N \delta(\vec{r}_j - \vec{r}) \delta(\vec{r}_k - \vec{r}_0). \quad (3)$$

It is crucial to analyze second order correlations, since the GS density only reveals radially symmetric contributions. Fig. 3 depicts $\hat{\rho}(\vec{r}, \vec{r}_0)$ for a selection of ground states with $N = 10$ particles. Generally, for small N , starting from the unstruc-

FIG. 3: Ground state density-density correlation functions $\hat{\rho}(\vec{r}, \vec{r}_0)$ for $N = 10$ dipolar fermions at L^z =(top) 45,80, (centered) 90,93, (bottom) 103,117 with $\vec{r}_0$ set to the maximum of the density, which occurs at the edge.



tured maximum-density-droplet at $L^z = N(N-1)/2$ (top left) with $\nu = 1$, a fraction of atoms starts to arrange itself in a “crystal” on the edge leaving a residual “Fermi sea” at the center (top right), until the correlations are homogeneously established at $L_0^z = 3N(N-1)/2$. For larger N , however, the energetically favorable number of edge atoms in the crystal changes “irregularly” with L^z (right column); the system seems to be “frustrated” in this respect. In this regime, the bulk has to reorganize itself accordingly, and novel fascinating states with a hole at the center of the bulk appear (centered and bottom left).

In order to understand the “magic” L^z numbers, Jain *et al.* [10] have proposed to model the interacting system in terms of an effective theory of non-interacting composite fermions (CF) for electrons in a quantum dot. In this ansatz, each fermion captures an even number of quantum fluxes, and the wave function reads

$$\Psi_{\text{CF}}(\{z_j\}) = \mathcal{N} \hat{\mathcal{P}}_{\text{LLL}} \left\{ \prod_{j < k}^N (z_j - z_k)^{2m} \Psi_{\text{Landau}} \right\}. \quad (4)$$

Here, m denotes the number of flux pairs, Ψ_{Landau} is an N -particle eigenfunction of $\mathcal{H}_{\text{Landau}}$, and $\hat{\mathcal{P}}_{\text{LLL}}$ is the LLL pro-

jector. Motivated by the success of CFs for fermions, Cooper and Wilkin have nicely adapted this scheme to bosonic contact interacting gases [14]. Following this idea, we compared the series of true GS for non-interacting composites and dipolar fermions. As long as there is no real bulk in the system, i. e., for $N < 7$, the ground state series nearly identically match with the predictions of the effective CF theory. Furthermore, overlap calculations for electrons and short-range interacting bosons [10, 14] suggest that similar results will hold for dipolar fermions. For bigger N , however, important deviations from CF theory, in particular in the intermediate regime, occur (see, for instance, the CF and true “magic” numbers in the table 5). This deficiency of the CF theory, already commented in Ref. [14], is more clearly related to the “frustration” effect and reorganization of the bulk in the present case. For $N = 10$, ground states with a density defect at the center are found at $L^z = 90, 95$ and 103 . They either have no ($L^z = 95$), or are hardly similar to the corresponding “magic” CF states. Instead, the analysis for the accessible range of systems strongly suggests that these states originate from “parent” states which are boosted by N quanta of angular momentum, e. g., $L^z = 80, 85$ and 93 for the above series. None of these parent states has its “magic” analogues in the CF model.

L^z , CF	45	55	63	69	77	83	90		97	103	111	117	125	135	(5)	
L^z , true	45	52	59	66	73	77	80	85	90	93	95	103	111	117		125


To a good approximation, the close connection between parent and boosted state (e.g. $L = 93$ and 103 respectively) can be understood as a quasi-hole excitation, which is analytically represented by

$$\Psi_{\text{qh}}(\{z_j\}) = \mathcal{N} \left\{ \prod_{j=1}^N (z_j - \zeta_0)^m \Psi_{L^z} \right\}, \quad (6)$$

where $\zeta_0 = 0$, and $m = 1$ for $\nu = 1/3$. Indeed, the above wave function proves to be a good approximation for the states with $L^z = L_{\text{parent}}^z + N$: even though the exact states do not reveal a true topological defect at the origin due to finite size, the density at the origin scales as $1/N$ and decreases from 0.14 to $\simeq 0.10$ as N varies from 7 to 11, while overlaps with the exact ground states grow from 0.6 to 0.7 despite the simultaneous significant increase of the relevant Hilbert space dimension.

We have also studied in detail low energy excitations of the dipolar Laughlin state at $\nu = 1/3$, which for $N \geq 10$ consists of a significant bulk, surrounded by a practically “melted crystal” at the edge (see Fig. 4). In order to relate our results to the ones obtained in the thermodynamical limit in Ref. [6], it is necessary to identify the finite-size quasi-hole excitations in the spectrum. To map out irrelevant candidates, it is reasonable to appropriately truncate the Hilbert space by a finite size

FIG. 4: Density-density correlation function $\hat{\rho}(\vec{r}, \vec{r}_0)$ of the Laughlin state for $N = 12$ dipolar fermions with $\vec{r}_0$ chosen at the maximum of the density, which occurs at the edge.



filling factor. We use the approach of Ref. [15], and fix ν by imposing the constraint $m_\nu \leq (N-1)/\nu$ on the maximum single particle angular momentum in the N -particle Fock basis. If a quasi-hole (quasi-particle) is nucleated, the filling factor is lowered (raised) by a correction of the order of $1/N$, i. e., $m_\nu \rightarrow m_\nu \pm 1$. This significantly reduces the basis dimension of the system and maps out irrelevant states. If the rotational frequency Ω is tuned to favor $\Psi_{1/3}^L$ as the true ground state, the lower lying excitations for $L_{1/3}^L < L^z < L_{1/3}^L + N$ turn out to be gapped by $\Delta\epsilon \simeq 1/N$. For short-range interacting


bosonic systems, these states have been partially identified as edge excitations that carry quanta of angular momentum [16]. To identify the excited states with a quasi-hole at the centre, the ground states at $L_{qh}^z = L_{1/3}^z + N$ are calculated with adapted $m_\nu + 1$. These states closely follow the lowest energy branch of edge excitations, and should be regarded as such, rather than quasi-holes. This is strongly supported by the fact that the substantial density dip for $N = 3$ continuously vanishes for increasing number of particles. Additional information about these edge excitations can be obtained by direct “naked-eye” inspection of Fig. 4. For the “standard” Laughlin state, the edge excitations are described by a hydrodynamical Luttinger liquid theory [17], and their amplitude decays as $\sin^{-3}(\phi)$, where ϕ is the “angular distance” from $\vec{r}_0$ along the edge; here the decay is much slower, which is yet another signature of the finite size effects.

The above considerations imply that the only reliable quantity that remains to estimate the quasi-hole energy in the large N limit is the neutral gap at fixed L^z . This gap remains more or less constant for smaller N s, and gradually increases as the number of bulk particles in the system and N grow. For the data of Ref. [6] ($M = 30$ a.m.u., $d = 0.5$ Debye, and a trap frequency of $2\pi \times 10^3$ Hz), the gap is of the order of percents of $2\hbar\omega_0$, i. e., it is substantial, but an order of magnitude less compared to the value estimated in [6] in the large N limit. Obviously, the lack of true bulk behavior in the investigated mesoscopic samples is responsible for this effect.

In the semiconductor fractional quantum Hall effect, Wigner crystals, i. e., specific charge-density waves, were discussed as competing ground states to Laughlin liquids. The interplay of quantum fluctuations with the interaction energy have proven crystalline Wigner order to be favorable for low enough fillings for electrons. This behavior intuitively changes in the case of dipolar particles as the interaction energy scales differently in the density of the particles. A very recent detailed analysis of this issue has confirmed the stability estimates for Laughlin and Wigner states of Ref. [6] for systems constituted of 50-200 particles [8]. Fehrmann et al. have proven that phonon excitations destabilize the Wigner crystal for fillings $\nu > 1/7$, which melts into a Laughlin liquid. In the microsystems discussed in this letter, this crossover is strongly supported. For fillings $\nu \leq 1/5$, density profiles significantly deviate from the Laughlin state and show a clear pattern of hexagonal order for $N = 6$ particles as depicted in Fig. 5 where the outer edge ring of the state is constituted of five particles. Even though speaking of a true “crystal” is certainly not applicable in such a system, signatures of crystalline order are clearly present.

Summarizing, we have studied in detail ground and excited states of quasi-2D ultracold rotating dipolar Fermi gases. By exact diagonalization methods, we studied systems up to 12 particles. We have identified novel kinds of strongly correlated states in the intermediate regime, i. e., pseudo-hole excitations of CF states. Calculation of the substantial gap in the excitation spectrum of the dipolar Laughlin state at $\nu = 1/3$ prove the accessibility of fractional quantum Hall states in

FIG. 5: Radial density $\rho(r)$ of Laughlin states (circles) and true dipolar ground states (triangles) for $N = 6$ at different filling factors. Apart from five particles which constitute the edge ring, localization of the sixth particle at the origin is clearly visible for $\nu \leq 1/5$.



these microsystems. At lower fillings, interactions favor crystalline order. Rotating dipolar gases are thus very suitable candidates to realize Laughlin-like and more exotic quantum liquids, as well as their crossover behavior to Wigner crystals.

We are indebted to W. Apel for valuable advice and help, and we thank D. Dagnino, M. Leduc, A. Riera, C. Salomon, L. Sanchez-Palencia, and G. V. Shlyapnikov for helpful discussions. We acknowledge support from the Deutsche Forschungsgemeinschaft (SPP1116 and SFB 407), ESF PESC “QUDEDIS”, EU IP “SCALA”, Spanish MEC (FIS2004-05639, FIS2005-04627, 2005SGR00343 and Consolider Ingenio 2010 “QOIT”).

-
- [1] M. Baranov, Ł. Dobrek, K. Góral, L. Santos, and M. Lewenstein, *Physica Scripta* **102**, 74 (2002).
 - [2] A. Griesmaier, J. Werner, S. Hensler, J. Stuhler, and T. Pfau, *Phys. Rev. Lett.* **94**, 160401 (2005).
 - [3] Special issue on *Ultracold Polar Molecules: Formation and Collisions*, *Eur. Phys. J. D* **31** (2004).
 - [4] N. R. Cooper, E. H. Rezayi, and S. H. Simon, *Phys. Rev. Lett.* **95**, 200402 (2005).
 - [5] S. Komineas and N. R. Cooper, cond-mat/0610702.
 - [6] M. A. Baranov, K. Osterloh, and M. Lewenstein, *Phys. Rev. Lett.* **94**, 070404 (2005).
 - [7] R. E. Prange and S. M. Girvin (editors), *The Quantum Hall Effect*, New York: Springer Verlag (1987).
 - [8] M. A. Baranov, H. Fehrmann, and M. Lewenstein, cond-mat/0612592.
 - [9] E. H. Rezayi, N. Read and N. R. Cooper, *Phys. Rev. Lett.* **95**, 160404 (2005).
 - [10] J. K. Jain, *Phys. Rev. Lett.* **63**, 199 (1989); J. K. Jain and T. Kawamura, *Europhys. Lett.* **29**, 321 (1995).
 - [11] This assumption is valid for systems subject to strong enough rotation such that the following conditions are fulfilled. The energy difference between LLs (modified by the trap and given by $\omega_+ = \sqrt{\omega_0^2 + \omega_c^2/4} + \frac{1}{2}\omega_c$ ($\hbar = 1$)) has to be much larger than the interaction energy, thermal interaction, and the single particle kinetic energy inside the LLL. The latter is given by

- $\omega_- = \sqrt{\omega_0^2 + \omega_c^2/4} - \frac{1}{2}\omega_c$; see A. G. Morris and D. L. Feder, cond-mat/0602037.
- [12] V. Fock, Z. Phys. **47**, 446 (1928);
C. G. Darwin, Proc. Camb. Phil. Soc. **27**, 86 (1931).
 - [13] E. R. Davidson, Computers in Physics **7**, 519 (1993).
 - [14] N. R. Cooper and N. K. Wilkin, Phys. Rev. B **60**, R16279 (1999).
 - [15] M. Kasner and W. Apel, Annalen Phys. **3**, 433 (1994).
 - [16] M. A. Cazalilla, Phys. Rev. A **67**, 063613 (2003).
 - [17] X.-G. Wen, *Quantum Field Theory of Many-Body Systems*, Oxford: Oxford University Press (2004).